

Louisiana Coastal Area (LCA), Louisiana

# **Ecosystem Restoration Study**

**July 2004**

**Draft**

**Appendix C – Hydrodynamic and Ecological Modeling**

**Page Intentionally Left Blank**

**LOUISIANA COASTAL AREA (LCA), LOUISIANA  
ECOSYSTEM RESTORATION STUDY**

**APPENDIX C  
HYDRODYNAMIC AND ECOLOGICAL MODELING**

**TABLE OF CONTENTS**

<b>FOREWARD.....</b>	<b>xix</b>
 <b>CHAPTER C.1 CONCEPTUAL ECOLOGICAL MODELS FOR PLANNING AND EVALUATION .....</b>	 <b>B-1</b>
1.1 INTRODUCTION .....	B-1
1.2 LCA STUDY AREA .....	B-2
1.3 CAUSAL MECHANISMS OF WETLAND LOSS IN COASTAL LOUISIANA.....	B-2
1.4 CONCEPTS OF RESTORATION SCIENCE.....	B-9
1.5 LCA ECOSYSTEM MODEL.....	B-15
 <b>CHAPTER C.2 FORMULATION OF THE LCA ECOSYSTEM MODEL.....</b>	 <b>B-18</b>
2.1 INTRODUCTION .....	B-18
2.2 LCA MODELING TEAM ORGANIZATION .....	B-18
2.3 LCA MODEL DEVELOPMENT.....	B-22
2.3.1 Existing Models .....	B-22
2.3.2 Modeling Tools.....	B-23
2.3.3 Simulation modeling.....	B-24
2.3.4 Desktop modeling.....	B-25
2.4 DATABASE FRAMEWORK DEVELOPMENT.....	B-26
2.4.1 Spatial Framework Development .....	B-26
2.4.2 Referencing of Key Spatial Datasets to the LCA Grid.....	B-27
2.4.3 Linking Channel Attributes to Wetland Regions.....	B-33

<b>CHAPTER C.3 HYDRODYNAMIC MODELS OF SUBPROVINCE 1 .....</b>	<b>B-36</b>
3.1 INTRODUCTION .....	B-36
3.2 MODEL OBJECTIVES.....	B-37
3.3 METHODS .....	B-37
3.4 MODEL DEVELOPMENT.....	B-39
3.5 BOUNDARY CONDITIONS .....	B-41
3.6 CALIBRATION AND VERIFICATION.....	B-45
3.7 RESULTS (BASE AND OPTIMIZED PLAN).....	B-47
3.8 DISCUSSION OF MODEL LIMITATIONS.....	B-52
 <b>CHAPTER C.4 HYDRODYNAMIC MODELS OF SUBPROVINCE 2 .....</b>	 <b>B-54</b>
4.1 IDEALIZED WINDS AND TIDES .....	B-54
4.1.1 Introduction.....	B-54
4.1.2 Objectives .....	B-55
4.1.3 Sensitivity Analyses.....	B-57
4.2 DYNAMIC WINDS AND TIDES .....	B-69
4.2.1 Introduction.....	B-69
4.2.2 Methods.....	B-70
4.3 RESULTS AND DISCUSSION .....	B-70
4.3.1 Barataria Bay Conditions in April and May, 1997 .....	B-70
4.3.2 Verification .....	B-73
4.3.3 Dynamic vs. Static Simulation.....	B-77
4.3.4 Testing the SRF Assumptions.....	B-78
4.4 MORE SENSITIVITY ANALYSES.....	B-80
4.4.1 Introduction.....	B-80
4.4.2 Methods.....	B-81
4.5 RESULTS AND DISCUSSION .....	B-82
4.5.1 Salinity Data.....	B-82
4.5.2 Model Runs.....	B-84
4.5.3 Analysis of Management Options.....	B-86
 <b>CHAPTER C.5 HYDRODYNAMIC MODELS OF SUBPROVINCE 3 .....</b>	 <b>B-89</b>
5.1 INTRODUCTION .....	B-89

5.2	METHODS .....	B-91
5.2.1	Hydrodynamic Module .....	B-91
5.3	RESULTS .....	B-92
5.3.1	Hydrodynamic Module .....	B-92
5.3.2	Near-Field Deltaic Land Building .....	B-93
5.3.3	Far-field Effects on Marsh Nourishment .....	B-94
5.4	ASSUMPTIONS AND LIMITATIONS .....	B-96
5.4.1	DATA VIGOR.....	B-96
5.4.2	SCIENTIFIC UNCERTAINTY.....	B-97
5.5	DISCUSSION .....	B-97
 <b>CHAPTER C.6 HYDRODYNAMIC MODELS OF SUBPROVINCE 4 .....</b>		<b>B-99</b>
6.1	INTRODUCTION .....	B-99
6.1.1	The Calcasieu-Sabine Basin .....	B-100
6.1.2	Approach.....	B-101
6.1.3	Model Setup.....	B-102
6.1.4	Boundary Conditions .....	B-103
6.1.5	Results (Base and Optimized Plan).....	B-103
6.1.6	Discussion .....	B-107
6.2	ROCKEFELLER WILDLIFE REFUGE SOUTH OF HWY 82.....	B-110
6.2.1	Introduction.....	B-110
6.2.2	Fresh Water Introduction Alternative .....	B-113
6.2.3	Project Description.....	B-114
6.2.4	Model Selection .....	B-114
6.2.5	Model Resolution.....	B-115
6.2.6	Data Collection & Review .....	B-115
6.2.7	Hydrologic Data Collection .....	B-116
6.2.8	Model Setup.....	B-118
6.2.9	Model Calibration .....	B-119
6.2.10	Model Validation .....	B-120
6.2.11	Evaluation of Model Performance .....	B-120
6.2.12	Discussion of Limitation and Capabilities of the Model .....	B-121
6.2.13	Initial Assessment of Project Features.....	B-123

6.2.14	Final Project Design Configurations.....	B-123
6.2.15	Salinity Analysis for the “Base Run” and “Final Design Run” .....	B-124
6.2.16	Discussion of Project Features and Impact on the Hydrology of the Region ..	B-126
6.2.17	Conclusions and Closing Remarks .....	B-126
6.2.18	Subprovince 4 Conclusions.....	B-126
<b>CHAPTER C.7 LCA Box Model Calculations.....</b>		<b>B-128</b>
7.1	BACKGROUND .....	B-128
7.2	INDEX MODELS (ORDER OF MAGNITUDE CHARACTERIZATION).....	B-128
7.2.1	Approach.....	B-128
7.2.2	Assumptions.....	B-128
7.2.3	Implementation .....	B-129
7.2.4	Output .....	B-129
7.3	BOX MODELS.....	B-130
7.3.1	Approach.....	B-130
7.3.2	Assumptions.....	B-130
7.4	CONCEPTUAL MODEL AND IMPLEMENTATION .....	B-131
7.4.1	Inputs from Other Modules.....	B-131
7.4.2	Outputs.....	B-131
<b>CHAPTER C.8 WETLAND NOURISHMENT MODULE.....</b>		<b>B-135</b>
8.1	INTRODUCTION .....	B-135
8.2	RATIONALE OF ASSUMPTIONS.....	B-135
8.2.1	Elevation Component.....	B-135
8.2.2	Land-Change Component .....	B-139
8.2.3	Land building.....	B-139
8.2.4	Nourishment Component.....	B-140
8.2.5	Spatial and Temporal Scales.....	B-143
8.3	DESKTOP ALGORITHMS .....	B-143
8.3.1	Land Building .....	B-143
8.3.2	Nourishment.....	B-144
8.3.3	Salinity Reduction Effect.....	B-145
8.3.4	Marsh Creation.....	B-145

8.4 RESULTS 146

**CHAPTER C.9 HABITAT SWITCHING MODULE.....B-148**

9.1	INTRODUCTION .....	B-148
9.2	RATIONALE OF ASSUMPTIONS AND INTERACTIONS .....	B-148
9.2.1	Habitat Switching Component.....	B-148
9.2.2	Habitat Productivity Component .....	B-152
9.3	SPATIAL AND TEMPORAL SCALES .....	B-159
9.4	DESKTOP ALGORITHMS .....	B-159
9.4.1	Habitat Switching Component.....	B-159
9.4.2	Habitat Productivity Component .....	B-160
9.5	RESULTS .....	B-162

**CHAPTER C.10 HABITAT USE MODULE.....B-166**

10.1	INTRODUCTION .....	B-166
10.2	LIMITATIONS OF THE MODULE.....	B-167
10.3	FISH AND SHELLFISH MODELS.....	B-168
10.3.1	General Methods.....	B-168
10.3.2	Juvenile Gulf Menhaden .....	B-170
10.3.3	Juvenile Spotted Seatrout.....	B-172
10.3.4	Juvenile Atlantic Croaker .....	B-174
10.3.5	Largemouth Bass .....	B-175
10.3.6	Brown Shrimp.....	B-177
10.3.7	White Shrimp.....	B-178
10.3.8	Oyster.....	B-179
10.4	WILDLIFE MODELS .....	B-180
10.4.1	General Methods.....	B-180
10.4.2	American Alligator .....	B-182
10.4.3	Dabbling Ducks .....	B-183
10.4.4	Mink.....	B-184
10.4.5	Muskrat .....	B-184
10.4.6	Otter .....	B-185
10.5	IMPLEMENTATION.....	B-186

10.6	RESULTS .....	B-186
<b>CHAPTER C.11 WATER QUALITY MODULE .....</b>		<b>B-189</b>
11.1	INTRODUCTION .....	B-189
11.2	APPROACH .....	B-190
11.3	METHODS .....	B-190
11.4	MODELS 195	
11.4.1	Inorganic N and Total Nitrogen Removal .....	B-195
11.4.2	Freshwater Flushing and Depth and Elevation Variables.....	B-199
11.4.3	Subprovince Total Percentage of N Removal.....	B-201
11.5	RESULTS 202	
<b>CHAPTER C.12 LCA BENEFITS ASSESSMENT PROTOCOLS .....</b>		<b>B-207</b>
12.1	INTRODUCTION .....	B-207
12.2	ECOSYSTEM OBJECTIVES .....	B-208
12.3	BENEFITS PROTOCOLS.....	B-208
12.4	SPATIAL AND TEMPORAL SCALES .....	B-209
12.5	BENEFITS PROTOCOL #1 (B1) .....	B-209
12.6	BENEFITS PROTOCOL #2 (B2) .....	B-210
12.6.1	Quality of Habitat .....	B-210
12.6.2	Quantity of Land.....	B-210
12.6.3	Nitrogen removal .....	B-211
12.6.4	Calculating B2 .....	B-211
12.7	BENEFITS PROTOCOL #3 (B3) .....	B-212
12.8	BENEFITS PROTOCOL #4 (B4) .....	B-212
12.9	BENEFITS PROTOCOL #5 (B5) .....	B-212
12.10	BENEFITS PROTOCOL #6 (B6) .....	B-214
12.10.1	Individual Species and Species Groupings.....	B-215
12.10.2	Habitat Types .....	B-215
12.11	LIMITATIONS .....	B-216
12.12	ACKNOWLEDGEMENTS.....	B-218

<b>CHAPTER C.13 MODEL UNCERTAINTY AND LIMITATIONS .....</b>	<b>B-219</b>
13.1 INTRODUCTION .....	B-219
13.2 UNCERTAINTY ANALYSIS, CONCEPTS, AND TERMINOLOGY .....	B-220
13.3 UNCERTAINTY ANALYSIS TECHNIQUES .....	B-222
13.3.1 Analytical methods .....	B-222
13.3.2 Random Sampling: Monte Carlo and Latin Hypercube methods .....	B-223
13.3.3 Bayesian approaches .....	B-223
13.3.4 Generalized Likelihood Uncertainty Estimation (GLUE) METHOD .....	B-224
13.4 SOURCES OF UNCERTAINTY IN THE LCA ECOSYSTEM MODEL .....	B-224
13.4.1 Sources of uncertainty in the Water Quality Module. ....	B-224
13.4.2 Sources of uncertainty in the Land change module. ....	B-226
13.4.3 Sources of uncertainty in the Habitat-switching module .....	B-230
13.4.4 Sources of uncertainty in the Habitat-use module .....	B-232
13.4.5 Uncertainty in Hydrodynamic models .....	B-236
13.4.6 Propagation of Uncertainties in the LCA Modules.....	B-241
13.5 VALIDATION OF THE CLEAR MODULES .....	B-244
13.6 UNCERTAINTY IN PERFORMANCE MEASURES .....	B-244
13.7 RECOMMENDATIONS ON FUTURE UNCERTAINTY ANALYSES FOR THE LCA ECOSYSTEM MODEL.....	B-245
<b>CHAPTER C.14 MODEL EVALUATIONS BASED ON SIMULATIONS OF A VIRTUAL BASIN   246</b>	
14.1 INTRODUCTION .....	B-246
14.2 SENSITIVITY ANALYSIS .....	B-246
14.3 VERIFICATION.....	B-249
14.4 VIRTUAL BASIN .....	B-252
14.5 REFINEMENTS .....	B-254
<b>CHAPTER C.15 MODELING AND MONITORING .....</b>	<b>B-256</b>
15.1 INTRODUCTION .....	B-256
15.2 MONITORING AND MODELING INTEGRATION.....	B-257

<b>CHAPTER C.16 LITERATURE REVIEW OF AVAILABLE DATA AND PREVIOUS STUDIES IN REGION 4: A BASIS FOR WATER AND SEDIMENT BUDGETS</b>	<b>260</b>
16.1 ABSTRACT/SUMMARY .....	B-260
16.2 BACKGROUND .....	B-262
16.3 AVAILABLE DATA.....	B-263
16.4 PREVIOUS MODELING EFFORTS WITHIN SUB-PROVINCE 4.....	B-265
16.5 DATA GAPS .....	B-266
16.6 MODELING AND ANALYSIS.....	B-266
16.6.1 Large Scale Hydrodynamic, Salinity, and Sediment Model.....	B-266
16.6.2 Sub-Province wide Hydrodynamic, Salinity, and Sediment Model: .....	B-266
16.6.3 WATER, SEDIMENT, AND SALINITY BUDGET ANALYSES.....	B-266
<b>LITERATURE CITED .....</b>	<b>B-268</b>

## List of Figures

Figure C.1-1	LCA Study Area and Subprovinces.....	B-2
Figure C.1 2	Seven Delta Lobes of the Mississippi River Delta .....	B-3
Figure C.1 3	Model of the Evolution of a Barrier Island System in the Mississippi River Deltaic Plain .....	B-4
Figure C.1 4	Conceptual Model of the Delta Cycle Depicting the Growth and Decay of a Delta Lobe .....	B-5
Figure C.1 5	Linkages in Land-Use Activities (including coastal restoration) .....	B-8
Figure C.1 6	Changes in Ecosystem Trajectories .....	B-10
Figure C.1 7	Goals of the LCA Ecosystem Restoration Plan Relative to Present Assumptions of Land Loss.....	B-11
Figure C.1 8	Conceptual Diagram Describing the Coastal Processes Wetland Ecosystem development.....	B-12
Figure C.1-9	Production Envelopes of Net Primary Productivity in response to the relative values of resource gradient, regulator gradient, and hydroperiod. ....	B-14
Figure C.1 10	16	
Figure C.2-1	Linkage of Different Modules used in Desktop and Simulation Models .....	B-19
Figure C.2-2.	Hybrid of desktop and simulation modeling tools for benefit evaluation. ....	B-24
Figure C.2-3	Schematic of how information was processed in the LCA Ecosystem Model using the GIS data base. ....	B-27
Figure C.2-4	Information used to generate the LCA desktop model habitat data base. ....	B-31
Figure C.2-5	Construction of spatially explicit landscape with over 38,000 one square kilometer (250 acres) cells. ....	B-32
Figure C.2-6	Boxes used to summarize information from hydrodynamic simulation models.	B-33
Figure C.2-7	Example of how information from the hydrologic simulations, at specific nodes in the channels of the coastal waters, is transferred to coastal wetlands at 1 km <sup>2</sup> resolution.....	B-34
Figure C.3 1	Example of Box Distribution and Boundaries in Subprovince 1.....	B-36
Figure C.3 2	Location of Subprovince 1 Study Area, Including Estuary Subdivisions .....	B-37
Figure C.3 3	Model Computational Domain and Bathymetry.....	B-40
Figure C.3 4	Tributary flows used in POM .....	B-41
Figure C.3 5	Diversion Hydrograph for the Base Alternative .....	B-42
Figure C.3 6	Diversion Hydrograph for the Reduce alternative .....	B-43
Figure C.3 7	Diversion Hydrograph for the Maintain Alternative .....	B-44

Figure C.3 8	Diversion Hydrograph for the Enhanced Alternative .....	B-44
Figure C.3 9	Observed and Modeled Water Surface Elevation at the Mid lake Station in Lake Pontchartrain .....	B-46
Figure C.3 10	Surface Salinity Distribution for Mid-April from the Future Without Scenario .....	B-47
Figure C.3 11	Surface Salinity Distribution for Mid-April from the Reduce Alternative .....	B-48
Figure C.3 12	Surface Salinity Distribution for Mid-April from the Maintain Alternative .....	B-48
Figure C.3 13	Surface Salinity Distribution for Mid-April from the Enhanced Alternative .....	B-49
Figure C.3 14	Monthly Mean Salinity for Future Without Alternative, Subprovince 1 .....	B-49
Figure C.3 15	Monthly Mean Salinity for the Reduced Alternative, Subprovince 1 .....	B-50
Figure C.3 16	Monthly Mean Salinity for the Maintain Alternative, Subprovince 1 .....	B-51
Figure C.3 17	Monthly Mean Salinity for the Enhanced Alternative, Subprovince 1 .....	B-52
Figure C.4 1	TABS-MD Model Domain as Developed by the USACE, New Orleans District.....	B-55
Figure C.4 2	Locations of 6 salinity output points in the lower Barataria Bay. Numbers inside the parenthesis indicate node numbers used in the “No-Marsh” model. ....	B-57
Figure C.4 3	Locations of cross-sections to compute inflow and outflow. Arrows and numbers indicate the flow directions and volume of flow in cfs respectively. ....	B-59
Figure C.4 4	Salinity plot with a 25 ppt constant Gulf boundary. ....	B-61
Figure C.4 5	Salinity plot with a 20 ppt constant Gulf boundary. ....	B-61
Figure C.4 6	Salinity plot for no wind condition. ....	B-63
Figure C.4 7	Salinity plot for 17.25 mph wind condition .....	B-63
Figure C.4 8	Salinity plot for variable wind condition. ....	B-64
Figure C.4 9	Salinity plot for 0 cfs diversion at Davis Pond. ....	B-64
Figure C.4 10	Salinity plot for 2500 cfs diversion at Davis Pond .....	B-66
Figure C.4 11	Salinity plot for 5000 cfs diversion at Davis Pond. ....	B-66
Figure C.4 12	Salinity plot for 7500 cfs diversion at Davis Pond .....	B-67
Figure C.4 13	Salinity Plot for 10600 cfs diversion at Davis Pond .....	B-67
Figure C.4 14	Salinity plot for 2500 cfs diversion at Davis Pond for 21 days, then off.....	B-68
Figure C.4 15	Salinity plot for 5000 cfs diversion at Davis Pond for 21 days, then off.....	B-68
Figure C.4 16	Grand Isle Tide for April and May 1997 .....	B-71
Figure C.4 17	Grand Isle Wind Speed for April and May 1997 .....	B-71
Figure C.4 18	Grand Isle Wind Direction for April and May 1997.....	B-72

Figure C.4 19	Location of Davis Pond SRC stations and LDWF Salinity stations 315 (Grand Terre) and 317 (St. Mary’s Point).....	B-72
Figure C.4 20	Observed salinity at St. Mary’s Point (317) and at Grand Terre (315) for April and May 1997 .....	B-73
Figure C.4 21	St. Mary’s Point Station 317 compared to model predictions for 500, 2,500 and 10,600 cfs Davis Pond diversion. Data from April and May 1997.....	B-74
Figure C.4 22	Grand Terre Station 315 compared to model predictions for 500, 2,500 and 10,600 cfs Davis Pond diversion. Data from April and May 1997.....	B-74
FigureC.4 23	Comparison of Predicted and Observed Salinities at Grand Terre for “no” Davis Pond discharge (actually 500 cfs) for April, 1997.....	B-77
Figure C.4-24	Salinity Plot 62 days with Grand Isle tide and wind. Davis Pond Diversion 2,500 cfs .....	B-79
Figure C.4 25	Salinity plot 62 days with Grand Isle tide and wind. Davis Pond Diversion 10,600 cfs .....	B-79
Figure C.4 26	Representative curve fit to USACE supplied salinity reduction factor nomographs. ....	B-80
Figure C.4 27	Location of Davis Pond SRC stations and LDWF Salinity stations 315 (Grand Terre) and 317 (St. Mary’s Point).....	B-81
Figure C.4 28	Stage at Venice and Barataria Salinities .....	B-84
Figure C.4 29	Effect of Boundary Salinity at Barataria Bay Water Way @ 15 ppt line .....	B-85
Figure C.4 30	Modeled Deviation of Bay Salinity at the 15 ppt Line Caused by 10 ppt Changes at the Gulf Boundary.....	B-86
Figure C.4 31	Hourly and Two Week Hindcast Salinities at the 15 ppt Line .....	B-88
FigureC.4 32	Effects of Gulf Salinities Changes at 5 ppt Line.....	B-88
Figure C.5 1	Location of Subprovince 3 with Superimposed Acadiana Basin Model Domain.....	B-90
Figure C.5 2	Detailed View of the Acadiana Basin Locations and Other Geographical Features of Interest .....	B-92
Figure C.6 1	The Chenier Plain of Coastal Louisiana .....	B-99
Figure C.6 2	General Layout of the Boundaries and Gauge Stations Used to Describe the Calcasieu-Sabine Basin.....	B-103
Figure C.6-3	Measured and Modeled Salinities at North Calcasieu Lake Near Hackberry .	B-104
Figure C.6-5	Comparison Between Measured and Modeled Salinity Values at Upper Sabine Lake at Platform “A” .....	B-105
Figure C.6 8	1999 Existing Water Level Results of North Calcasieu Lake Near Hackberry, LA .....	B-107
Figure C.6-9	View of March 1999 for North Calcasieu Lake near Hackberry, LA .....	B-108

Figure C.6 10	Salinities at North Calcasieu Lake near Hackberry, LA.....	B-108
Figure C.6-11	Typical Model Output Showing a Salinity Map at the Water Surface for the Calcasieu-Sabine Basin.....	B-109
Figure C.6-12	Project Location Map.....	B-110
Figure C.6 13	Base Map .....	B-111
Figure C.6 14	Existing Control Structures Within and Near the Project Area .....	B-113
Figure C.6 15	Basemap Showing Location of Continuous Recorders Used for the Model ...	B-116
Figure C.6-16	Salinity Model Results Compared to Field Data at ME16-03 .....	B-120
Figure C.6-17	Water Level Model Results Compared to Field Data at ME16-03.....	B-120
Figure C.6-18	Salinity Model Results Compared to Field Data at ME16-03 (Validation).....	B-122
Figure C.6-19	Water Level Model Results Compared to Field Data at ME16-03 (Validation).....	B-122
Figure C.7 1	Factors affecting aggregated salinity in each region.....	B-132
Figure C.7 2	Process of calibrating mass-balance models based on results of hydrodynamic models.....	B-133
Figure C.7-.3	Residence time for mixing can be inferred from the decay in concentration of a conservative tracer in the model .....	B-134
Figure C.8-1	Conceptual Model of Elevation Change Used in Existing Soil Formation Simulation Models .....	B-136
Figure C.8-2	Distribution of existing Cesium core data analyzed for this module.....	B-137
Figure C.8-3	Accretion rates by habitat and river influence. Error bars reflect one standard error. River influenced rates are based on four or fewer cores per marsh type.....	B-137
Figure C.8-4	Subsidence rates for the Louisiana coastal area (based on Kulp 2000).....	B-138
Figure C.8-5	Generalized order at which cells will be filled with sediments from the diversion. This order is determined based on the distance of the cell from the diversion.	B-140
Figure C.8-6	Land change pattern away from the mouth of the Atchafalaya Delta. Only the east side of the mouth was used, because of the effect of the Wax Lake Delta to the west. ....	B-141
Figure C.8-8	Distribution of Study Sites Used to Test Effect of Caernarvon Diversion on Land Change Rates .....	B-142
Figure C.8-9	Wetland area in Year 50 as a Result of Different Restoration Scenarios in Order of Increased Sediment Load.....	B-146
Figure C.8-10	Example of the Spatial Representation of Results from the Land Change Module .....	B-147
Figure C.9 1	Conceptual Model for the Habitat Switching Component.....	B-150

Figure C.9 2	Simplified Model for the Habitat Switching Component .....	B-150
Figure C.9-3	Conceptual Diagram Demonstrating Interaction Between Salinity and Inundation Restrictions of Brackish and Saline Marsh Vegetation .....	B-152
Figure C.9 4a	Effect of Salinity on Production of Swamp Forests as Derived from a Literature Review .....	B-154
Figure C.9-4b	Effect of Salinity on Production of Fresh Marsh with Saturated Soils as Derived from a Literature Review .....	B-154
Figure C.9-4c	Effect of Salinity on Production Intermediate Sagittaria Marsh as Derived from a Literature Review.....	B-155
Figure C.9.4d	Effect of Salinity on Production of Intermediate Spartina Marsh as Derived from a Literature Review.....	B-155
Figure C.9-4e	Effect of Salinity on Production of Brackish Marsh as Derived From a Literature Review .....	B-156
Figure 9.4f	Effect of Salinity on Production of Saline Marsh as Derived from a Literature Review .....	B-157
Figure C.9-5	Spatial Distribution of Average Annual Salinity in Subprovince 1 Under Restoration Scenario M02.....	B-164
Figure C.9-6	Spatial Distribution of Habitats in Subprovince 1 Under Restoration Scenario M02 .....	B-164
Figure C.9-7	Change in Habitat Composition Resulting from Different Restoration Scenarios .....	B-165
Figure C.9-8	Changes in Marsh Productivity Index Resulting from Different Restoration Scenarios .....	B-165
Figure C.10 1	Conceptual Model.....	B-169
Figure C.10 2	Restoration Scenario Effects on Habitat Suitability for Fish and Shellfish Species in Subprovince 1 .....	B-187
Figure C.10 3	Restoration Scenario Effects on Habitat Suitability for Wildlife Species .....	B-188
Figure C.11-1	Example of Box distribution and boundaries in subprovince 1 (Breton/Pontchartrain) used to evaluate water quality variables. ....	B-191
Figure C.11-.2	Initial LCA mapping categories and new categories defined to obtain the land water and volumes for the water quality module.....	B-192
Figure C.11-3	Flow chart of the water quality module to estimate suitability indexes (HSI) .....	B-193
Figure C.11-4	Chlorophyll a, productivity, and nitrogen removal suitability indexes .....	B-194

Figure C.11-5 Relationship Between Freshwater Residence Time and Fraction of N Exported .....	B-196
Figure C.11-5 Fraction of Upland Nitrogen Input Exported versus Freshwater Residence Time .....	B-196
Figure C.11-6 Nitrate-Nitrogen Removal by Mass and Concentration versus Nitrate-Nitrogen Loading .....	B-197
Figure C.11-7 Relationship between Percentage of TN Loading and Removed by Denitrification and Water Residence Time .....	B-197
Figure C.11-8 Scatter Plot Relating Annual Areal TN Loads to Annual Average Chlorophyll a Concentrations .....	B-198
Figure C.11-9 Relationship to Estimate Primary Production in the Water Column .....	B-198
Figure C.11 10Schematic showing changes in ‘wetland water’ base depth before and after inundation. ....	B-201
Figure C.11-11Conceptual diagram showing the overall calculation of N removal for each subsystem (boxes) within each subprovince. ....	B-202
Figure C.11 12Percentage of Total Nitrogen Removal for Each Scenario by Subprovince....	B-206
Figure C.13-1 Sources of types of uncertainty considered in current analysis.....	B-222
Figure C.13-2 Illustration of the Monte Carlo simulation method formodel uncertainty analyses .....	B-224
Figure C.13-3. Average total monthly sediment load and flow calculated for 10-year increments, showing the amount of interdecadal variation. ....	B-229
Figure C.13-4 Schematic diagram showing exchange of information and associated uncertainties across the LCA modules. ....	B-243
Figure C.14-1 Sensitivity of the Emergent Marsh Productivity Index to Salinity .....	B-248
Figure C.14-2 Sensitivity of Habitat use Model to Salinity and Wetland Area Criteria.....	B-249
Figure C.14-3 LCA Alternatives from Subprovince’s 1-3 Ranked by Total Discharge .....	B-250
Figure C.14 4 Correlation Between Discharge and Sediment Load in Three Subprovinces..	B-251
Figure C.14-5 LCA Alternatives from Subprovince’s 1-3 Represented by Discharge and Land Acreage in Year 50 .....	B-252
Figure C.14-6 Discharge Effect on Total Vegetative Production .....	B-252
Figure C.14-7 Relationship Between Land Acreage and Total Productivity Index for Year 50.....	B-253
Figure C.14-8 Discharge Effects on Cumulative HSI for High Salinity Species (Oyster, Brown Shrimp, Spotted Seatrout).....	B-254

Figure C.14-9 Hydrographs of the Virtual Basin Representing 1,000 cfs, 10,000 cfs and 100,000 cfs Average Annual Discharge into the Upper and Lower Basin.....	B-254
Figure C.14-10 Total Wetland Area at Year 50 in the Virtual Basin .....	B-255
Figure C.14-11 Total Wetland Area at Year 50 and Annual Sediment Load in the Virtual Basin .....	B-256
Figure C.15-1 Linkage of modeling and monitoring programs to develop adaptive anagement and assessment of restoration planning and evaluation.....	B-259
Figure C.15-2 Spatial Orientation of CRMS-Wetland Stations .....	B-260
Figure C.16-1: General map showing the boundaries of the Chenier Plain .....	B-261
Figure C.16-2: Sample of available bathymetric data in sub-province 4 at the time of writing this report. ....	B-264
Figure C.16-3: Sample of the continuous monitoring stations in sub-province 4. ....	B-265
Figure C.16-4: Sample of the historical discrete data in sub-province 4.....	B-265
Figure C.16-5: Additional historical discrete data in sub-province 4.....	B-266

## List of Tables

Table C.2-1	List of participants in the LCA Ecosystem Model program.....	B-20
Table C.2-1	Description of the alternatives simulated with numerical models.....	B-25
Table C.2-2	LCA Spatial Framework Parameters .....	B-27
Table C.2-2	Spatial Analysis Terms .....	B-28
Table C.2-3	Key LCA Spatial Datasets .....	B-28
Table C.2-4	2001 Linscombe Marsh Type Classes .....	B-30
Table C.2-5.	Number of 1 km <sup>2</sup> (250 acres) cells in the four subprovinces of the LCA Ecosystem Model. ....	B-32
Table C.3 1	Error analysis for POM in Lake Pontchartrain. ....	B-46
Table C.3.2	Observed and Modeled Currents in Lake Pontchartrain.....	B-46
Table C.4 1	Conservation of mass (flow) test for RMA2 (hydrodynamics) .....	B-58
Table C.4 2	Effect of Varying Boundary Salinity on Target Salinity (31 day average, all values in ppt).....	B-60
Table C.4 3	Effect of Varying Wind Speed on Target Salinity (South Wind, 31 day average, all values in ppt).....	B-62
Table C.4 4	Effect of Davis Pond Discharge on Target Salinity*.....	B-69
Table C.4 5	Salinity Reduction Associated with each 1,000 cfs of Davis Pond Discharge After 31 Days .....	B-69
Table C.4 6	Comparison of Predicted and Observed Salinities for April and May 1997 .....	B-73
Table C.4 7	Effect of Varying Davis Pond Discharge on Target Salinity with Jan & Feb 1997 Winds and Tides .....	B-75
Table C.4 8	Comparison of Monthly Salinities Observed and Predicted Using the "No Marsh" TABS and the SRF Method .....	B-76
Table C.4 9	Hourly Salinity Data at Target Locations: 12/8/ 2001 to 3/13/ 2002 .....	B-82
Table C.4 10	Target Salinity Ranges by Month at the 5 ppt Salinity Line.....	B-83
Table C.5 1	Annual Sediment Supply and Percent Retention in the WLO and LAR Deltas .....	B-93
Table C.5 2	Land Change Results of 70-Year ABM Runs.....	95
Table C.5 3	Observed and Predicted Effects of Discharge on Annualized Land Change Rates (%-y-1).....	96

Table C.5 4	ABM Prediction of the Influence of River Discharge on Land Gain and Loss Within the Model Domain (km <sup>2</sup> yr <sup>-1</sup> ) .....	B-97
Table C.6 1	List of Alternatives for the Calcasieu-Sabine Basin .....	B-101
Table C.6 2	Summary of Error Analysis .....	B-109
Table C.6 3	Description of Existing Water Control Structures .....	B-112
Table C.6 4	Availability of Data for the ME-16 Modeling of Hwy. 82 Freshwater Introduction Project .....	B-118
Table C.6-5	Quantitative Assessment of Model Results .....	B-121
Table C.8 1	Land loss rate categories based on land area changes between 1978 and 1990.....	B-139
Table C.8 2	Change in Percentage Land Area Per Year with Distance From the Diversion Point .....	B-145
Table C.9-1	Restrictions in Salinity and Inundation for the Major Habitat Types1 .....	B-159
Table C.9-2	Salinity Assumed to Have 100% Production and Regression Coefficients for the Different Habitat Types .....	B-161
Table C.9-3	Production Values Compiled from the Literature for Each Habitat Type .....	B-167
Table C.9-4	Percent of Annual Primary Productivity by Season and Habitat Type.....	B-168
Table C.10 1	Factors and Variables Available for the Habitat Suitability Analysis for each of the 0.3 mi <sup>2</sup> (1km <sup>2</sup> ) Cells .....	B-167
Table C.11-1	Flushing Times in Several Coastal Systems .....	B-200
Table C.11 1	Suitability Indices for Subprovince 1 (Mississippi East (Breton) .....	B-203
Table C.11 2	Suitability indexes Indices for Subprovince 1 West (Pontchartrain).....	B-204
Table C.11 3	Data Summary .....	B-205
Table C.12 1	Summary Description of LCA Benefits Protocols.....	B-208
Table C.12 2	Species Included in Benefit and Variable Designations .....	B-212
Table C.12 3	Variable Weights (%) for Each Model .....	B-214
Table C.12-4	Species Grouping used in Benefits Protocol #6.....	B-215
Table C.12 5	Summary of Benefits Protocol Calculations.....	B-216
Table C.13-1.	Overview of water quality model uncertainty. ....	B-225
Table C.13-2.	Definition and values of parameters used in the Water Quality Module.....	B-227
Table C.13-3.	Overview of land change model uncertainty. ....	B-228
Table C.13-4.	Definition and values of parameters used in land change model .....	B-232
Table C.13-5.	Overview of Habitat Switching Model Uncertainty .....	B-233
Table C.13-6.	Overview of Habitat Use (Wildlife) Model Uncertainty .....	B-237

Table C.13-7 Overview of Habitat Use (Fisheries) Model Uncertainty .....	B-237
Table C.13-8 Overview of Hydrodynamic Model Uncertainty .....	B-238
Table C.13-9. Definition and values of parameters used in the Princeton Ocean Model .....	B-240
Table C.13-10. Definition and values of parameters used in the Acadiana Basin Model .....	B-241
Table C.13-11 Module variables and interactions among modules.....	B-244

## Foreword

This Louisiana Ecosystem Study (LCA) Report Appendix entitled, “Hydrodynamic and Ecological Modeling” is designed to provide an understanding of the basis on which initial subprovince wide model efforts were conducted. These efforts provided the LCA Project Delivery Team (PDT) with tools to evaluate various proposed restoration measures. The modeling effort summarized here was conducted during a period between August 2002 and September 2003. This modeling effort developed tools used to select the seven original coastwide frameworks developed through phase five of the LCA plan formulation (see Main Report).

The models represent a landmark achievement in collating the extensive scientific and technical knowledge currently available. They were created for the purposes of formulating and evaluating subprovince-level alternative plans (which were comprised of multiple project features) for ecosystem restoration. Although the lack of extensive data sets constrains the models' ability to accurately predict individual project benefits over 50 years, the conceptual frameworks developed for the models are sound and represent the best available scientific understanding of ecosystem function. Even though precise project benefits over 50 years could not be predicted, resource managers were able to compare the relative effectiveness of alternative subprovince plans. They could also distinguish between these plans with reasonable certainty that the most cost-effective and ecologically beneficial alternatives were being considered. Further, these models represent the most objective and powerful predictive tool at the subprovince scale available to resource managers at this time.

Output provided by these models assisted the PDT in the determination of 7 cost-efficient alternative plans from which cost-efficient and ecologically meaningful projects could be chosen. The projects that comprised each of the 7 plans were considered in the development of the near-term plan for the Louisiana Coastal Area (LCA) Study. Upon completion, only 13 of the 79 project features considered passed through the selection process for inclusion in the near-term plan. Of those 13 projects, only 5 are being recommended for programmatic authority for construction. These 5 have been selected because they have significant value in addressing critical ecological needs, they have some level of planning and design already completed, and because they will utilize technology which has already been proven to be cost-efficient and ecologically beneficial by similar projects implemented under other Federal and State programs. This is also consistency with the Adaptive Environmental and Assessment and Management (AEAM) process, described elsewhere in this report, wherein the design of future actions is built upon lessons learned regarding the efficacy of past restoration actions.

Uncertainty is inherent in ecosystems, and is therefore unavoidable when managing large-scale ecological systems. Thus, assumptions must be made when creating predictive ecological tools. As acknowledged above, the lack of extensive data sets for all parameters being considered creates further uncertainty in the models' ability to accurately predict benefits

over the 50-year project life. Acknowledging and identifying these and other uncertainties is critical for the most appropriate utilization of output. It is the consensus of the scientists who created these models that the outputs are a sound basis for decision making at the subprovince scale.

As the LCA Program proceeds, these subprovince-level models will continue to be developed and – where possible – uncertainty will be reduced through the Science and Technology Program. This is an integral step in the LCA AEAM program, and it allows for large-scale ecosystem restoration to proceed even as researchers work to reduce those uncertainties.

## CHAPTER C.1

# CONCEPTUAL ECOLOGICAL MODELS FOR PLANNING AND EVALUATION

Robert R. Twilley <sup>1</sup>

<sup>1</sup> *Center for Ecology and Environmental Technology, University of Louisiana at Lafayette*

### 1.1 Introduction

The proposed LCA Ecosystem Restoration plan establishes a framework or blueprint for solution of the Louisiana coastal problems and opportunities for wetland rehabilitation over the near term of 10 years. The near-term course of action restoration opportunities capitalize on the set of coastal framework features developed by the LCA Study team's ecological modeling efforts.

This appendix provides a summary of the conceptual ecological modeling process utilized to support the planning and evaluation processes of the LCA Ecosystem Restoration Plan. Further, this appendix focuses on the following modeling tasks to support the evaluation of proposed coastwide restoration frameworks: 1) Development of 'Conceptual Ecological Models' used to integrate ecological needs and opportunities with engineering designs that provide the most benefit to coastal Louisiana ecosystems; 2) Use rates of wetland loss to describe the most likely future without scenario for variety of ecosystem attributes; 3) Establishment of broad ecosystem responses to restoration alternatives based on processes associated with succession of geomorphic and ecological systems; 4) Assessed water and sediment needs from the Mississippi and Atchafalaya rivers, and other sources, to establish site criteria necessary to obtain restoration goals within the Mississippi River Delta; 5) Development of ecological benefits assessment protocols that associate large-scale geomorphic and hydrologic processes that would lead to rehabilitation of the Mississippi River deltaic and Chenier Plains.


The Applied Science Strategy approach outlined in the Comprehensive Everglades Restoration Plan (Appendix D Attachment A, pages 1-90 of the Central and Southern Florida Project Comprehensive Review Study) was modified for the specific site conditions of coastal Louisiana to provide a level of detail required for engineering and environmental aspects of plan formulation and justification of a LCA coastwide study. The Applied Science Strategy described focuses on Conceptual Ecological Models of both the natural system degradation and needs for ecosystem rehabilitation to make sure there are clear statements of problems, needs and opportunity in the planning phases. The initial step of this Conceptual Model was to define disturbances, sources of ecosystem stress, and development of desired ecosystem response. These assumptions are based on clear causal linkages between disturbances and ecological effects. Second, the model describes desired ecological endpoints or restoration response based on the principles of sustainable ecosystem processes in deltaic environmental settings (self-design). These responses require an understanding of present ecosystem state, desired endpoints, and necessary site conditions to obtain specific endpoints.

The focus of the efforts in the LCA Ecosystem Model was to add a level of detail in the causal linkages, rationale on desired site conditions, and engineering requirements to reach

landscape and ecosystem endpoints that are the goal of this ecosystem restoration plan. Chapter D.2 of this appendix describes the formulation of a LCA Ecosystem Model that uses conceptual frameworks and ecosystem objectives to design an effective tool in coastal restoration science. Chapters D3 through D6 discuss the use of hydrodynamic models utilized for each subprovince to provide input for the LCA conceptual ecological model. Chapters 7 discusses the LCA box model with specific parameter models discussed in Chapters 8-11. The protocols for assessment of benefits is presented in Chapter 12. Chapters 13-15 discuss model evaluation, limitations, uncertainties and future requirements for refinement.

## 1.2 LCA Study Area

The study area, which includes Louisiana's coastal area from Mississippi to Texas, is made-up of two wetland-dominated ecosystems, the Deltaic Plain of the Mississippi River and the closely linked Chenier Plain, both of which are influenced by the Mississippi River. For planning purposes, the study area was divided into four subprovinces, with the Deltaic Plain comprising Subprovinces 1, 2, and most of 3, and the Chenier Plain comprising the western part of Subprovince 3 and Subprovince 4 (**Figure C.1-1**). The Mississippi River Deltaic and Chenier Plains consists of diverse geomorphological basins with distinct vegetation zones and patterns of development.




**Figure C.1-1 LCA Study Area and Subprovinces**

## 1.3 Causal Mechanisms of Wetland Loss in Coastal Louisiana

Deposition of sediments by the Mississippi River led to the formation of the present Mississippi delta and associated chenier plain, which is composed of more than 9.9 million acres

(4.0 million ha) of wetlands, lakes and bays (Roberts 1977). Discharge of fresh water and sediment from the Mississippi River has changed course over the last 7000 years (Figure C.1-2), resulting in the formation of two distinct provinces along the coast: a deltaic plain to the east and Chenier Plain to the west (Boesch *et al.* 1994). The delta is ecologically diverse and productive and economic activities depend on the productivity of this natural resource. The Mississippi Deltaic Plain is characterized by high riverine input, shallow bays, vast wetlands, in a warm temperate, low energy coastline. The Mississippi River Deltaic Plain has the eighth highest annual mean freshwater discharge in the world, 640,000 ft<sup>3</sup>/s (18,000 m<sup>3</sup>/s), which causes extreme spatial and temporal variation in distribution of particulate and dissolved materials within the coastal waters. The broad continental shelf and prevailing winds tend to isolate the land margin from open ocean processes such as upwelling of slope waters. Daily tidal amplitudes are small, averaging only 0.98 ft (30 cm), but water level fluctuations over 3.3 feet (1 m) can occur during frontal passage. On a longer time scale the relative rise in sea level in this region is in excess of 0.033 ft/yr (1 cm/yr) due primarily to a high rate of regional subsidence. In addition, the hydrodynamics of the region are directly influenced aperiodically by hurricane surges that occur about once every 5 yrs. Thus significant water level changes in the coastal margin of the Mississippi Deltaic Plain range from daily to geologic time scales.




Source: Gosselink 1998; original modified from Kolb and Van Lopik 1958

**Figure C.1-2 Seven Delta Lobes of the Mississippi River Delta**

The delta cycle is fundamental to understanding the succession of geomorphic and ecologic features of this coastal landscape (Figure C.1-3). Transgressive sequences at the province and basin scales of coastal Louisiana govern smaller scale successional changes at the habitat scale of the marsh. The proximity of fluvial processes to marshes shift as distributaries of the Mississippi River migrate along the coast, changing the distribution of sediment, nutrients, and salt that control the type of habitat that colonizes the emergent zones of the basin. Thus there are continued changes not only from emergent to open water as part of the transgressive sequences, but the community composition of the emergent lands changes among fresh water, intermediate, brackish, and salt marsh vegetation (Figure C.1-4). As fluvial processes decrease,

there is a lack of fresh water discharge to control sea water encroachment, causing salt and brackish marshes to migrate landward, either replacing fresh water marshes or converting marshes to open water (Figure C.1-4). During active delta formation, such as observed in the Atchafalaya River basin, there is a migration of fresh water and intermediate vegetation toward the coast as salinity regimes decrease in the coastal zone (Madden *et al.* 1988). Processes at all three spatial scales including province, basin and habitat levels are coupled to produce a spatial mosaic of changes in wetland cover and composition that form very complex and dynamic patterns of coastal wetland succession. The result of these processes across the Mississippi River Deltaic Plain is 6,200,000 acres (2,500,000 ha) of marshes that account for 60% of the coastal wetlands in the lower 48 states (Turner and Gosselink 1975). These patterns of coastal processes have to be incorporated in any perspective of coastal restoration and rehabilitation.




Source: Gosselink 1998; Original from Penland *et al.* 1988

**Figure C.1-3 Model of the Evolution of a Barrier Island System in the Mississippi River Deltaic Plain**

The Mississippi River actually formed two distinct geomorphic regions of coastal Louisiana over the last 7000 years – the Deltaic Plain and the Chenier Plain. The Deltaic Plain formed in the central and southeastern portions of the coast (Fisk and McFarlan 1954), and the Chenier Plain in the southwestern part of the state (Gould and McFarlan 1959; Penland and Suter 1989). The hydrology and landscape formation of marshes in these two coastal regions are distinctly different. Subsidence in the deltaic region of coastal Louisiana averages 0.036 ft/yr (1 cm/yr) compared to 0.019ft/yr (0.57 cm/yr) in the Chenier Plain region (Penland and Ramsey 1990). These differences in regional subsidence rates are associated with the erosion of Pleistocene surfaces by the Mississippi River in the Deltaic Plain followed by deposits of fine silts to depths of over 656 ft (200 m) there (Penland and Suter 1989). In the Chenier Plain, much less erosion occurred because the river never flowed directly through the region; therefore depths to Pleistocene surfaces are only 49 ft (15 m) (Nichols 1959). In the Mississippi River Deltaic Plain, wetlands initially form as freshwater marshes at the mouths of active distributaries and

convert to saline marshes as the delta lobe cycle progresses toward the degradation phase (Coleman 1988). In the Chenier Plain, wetlands initially form as saline marshes in the Gulf of Mexico and convert to freshwater marshes as new marshes and chenier isolate them from the Gulf of Mexico.



Source: Gosselink 1998, modified from Gagliano and Van Beek 1975; Neill and Deegan 1986

**Figure C.1-4 Conceptual Model of the Delta Cycle Depicting the Growth and Decay of a Delta Lobe**

The Mississippi River Deltaic and Chenier Plains consists of diverse geomorphological basins with distinct vegetation zones and patterns of development. Within each of these geomorphological basins are ecological habitats that can be distinguished by the adaptation of

plants to soil fertility, relative water levels, and salinity (Buresh *et al.* 1980, DeLaune *et al.* 1989). Marshes in coastal regions that differ in geomorphology and nutrient (fresh water and sediments) loading have different plant strategies to resource availability and abiotic stressors (Hopkinson and Schubauer 1984; White and Howes 1994c), and thus different patterns of marsh stability. Natural shifts in proximity to fluvial inputs will change the relative loadings of N and P (and Fe), as well as salinity, and this should cause shifts in relative production and decomposition of organic matter in marsh wetlands (Day *et al.* 1995). The conceptual model of the delta lobe cycle predicts that diversity and possibly productivity actually peaks following the peak in land mass formation, with slight increase in salinity during the early phases and delta degradation.

The biological productivity of this coastal landscape is linked to the extensive diversity of coastal habitats in this geographically distinct central Gulf Coast region. The biological diversity and productivity of the Mississippi River delta includes the largest wetland landscape and most productive near shore fishery and migratory bird habitat in the contiguous United States. Much of this biological diversity and productivity of these higher trophic levels is associated with the delta cycle and corresponding distribution of land mass and salinity regimes. Thus the succession of wetland vegetation and habitat development influences the relative dominance of marine and terrestrial fauna. One of the unique features of this broad landscape from the delta to the chenier plain is the seasonal use of these habitats by fauna during critical stages of their life cycle.


Coastal Louisiana has the highest rate of coastal wetland loss in the nation, reaching a peak of 0.86% per year 41.7 mile<sup>2</sup>/yr (108 km<sup>2</sup>/yr) in the 1970's (Gagliano *et al.* 1981; Turner and Cahoon 1987; Barras *et al.*, Appendix G). Although the rate has declined since the 1980's (Britsch and Dunbar 1993), over one-third of the wetlands present in 1930s have been lost, equivalent to an area equal the size of Rhode Island (Boesch *et al.* 1994). This has resulted in the conversion of vegetated areas to open water decreasing the wetland:water area ratio. A large proportion of this loss occurs as conversion of interior marshes of all types (salt, brackish, and fresh) to open water (Gagliano *et al.* 1981; Turner and Cahoon 1987; Penland *et al.* 1996). Hence, wetland loss is related not only to erosion of the marsh edge but also factors contributing to submergence of interior wetlands. Much of this wetland loss is associated with a high rate of regional subsidence and erosion characteristic of degrading deltas (Figure C.1-4). At the mouth of the Atchafalaya River, however, a wetland system representing the early progradational stages of delta formation is evolving (van Heerden 1983). Here the levels of sediment discharge compensate for the relative increase in water levels due to subsidence and sea level rise (Day *et al.* 1995). This huge landscape of wetlands and coastal bays fluctuates in total area depending on a balance between the progradational processes of active delta formation, and degradational processes during abandonment that lead to natural and unnatural wetland loss (Figure C.1-4).

There are several factors in the environmental setting of coastal Louisiana which contribute to an inability of wetlands to maintain surface elevation leading to marsh instability (Figure C.1-5): (1) a high rate of regional subsidence (Penland and Ramsey 1990); (2) a reduced sediment load in the Mississippi River (Kesel 1988); (3) elimination of spring overbank flooding of the Mississippi River and direct delivery of river sediment to the marshes (Templett and Meyer-Arendt 1988; Day *et al.* 1997); and (4) extensive landscape and hydrologic alterations from human activities, including energy related activities and navigation channels (Turner and Cahoon 1987). Determining which process is most important at controlling elevation (*e.g.*, mineral vs. organic matter accumulation; waterlogging vs. salinity stress) is important in

developing a rehabilitation program. In a comprehensive evaluation of the wetland loss problem in Louisiana, a panel of expert coastal scientists recommended that fundamental emphasis should focus “...on processes of soil formation, including the importance of mineral sediments in different types of wetlands, geochemical conditions affecting organic matter incorporation into sediments, and historical changes in soil characteristics.” (Boesch *et al.* 1994).

Levees along the Mississippi River were initially constructed in the late 1700s to prevent waters during high river stage from flooding homes and agricultural fields, causing death and economic ruin to families all along the river. Several catastrophic breaches occurred as a result of record flood events along the river as recently as the early 1900s, each resulting in repairs to the levee system improving its capacity to prevent flooding. Thus, with the exception of short periods of time up to the early 1900s, the levees have been in place for over 100 years. However, extremely high rates of land loss along the coastal landscape of Louisiana during the late 1900s brought attention to the potential impacts of limited river flooding in hydrologic basins. Land loss was attributed to sea level rise, land subsidence, and the lack of river sediments being delivered to the marshes. Coastal marshes in deltaic environments are highly dependent on land building by soil formation that is enhanced by periodic delivery of rich new sediments from river floodwaters, as described above. The Mississippi River levee prevented sediment deposition from occurring in deltaic marshes, prompting managers to find ways of restoring river flow to impacted coastal areas.

Land loss in the Chenier Plain was been attributed largely to erosion of shoreline resulting in the loss of habitat; and salt water intrusion that has converted many marshes to open water. In addition, there is evidence that increased water logging in some regions may contribute to wetland instability. It is important that casual mechanisms of wetland loss be specific to the deltaic and chenier plain in any recommendation of coastal rehabilitation.



**Figure C.1-5 Linkages in Land-Use Activities (including coastal restoration)**

Marsh stability is critically linked to the relationship between marsh elevation and soil formation. Processes influencing elevation include mineral sedimentation, organic matter production, organic matter decomposition, and subsidence (Figure C.1-5). When sea level rises faster than marsh elevation, an aggradation deficit develops resulting in the eventual submergence of the marsh. The productivity and health of the vegetation are central to maintaining this balance because wetland soils in coastal Louisiana depend on marsh vegetation to create enough soil organic matter not only to create new soil, but to also offset soil organic matter decomposition (Callaway *et al.* 1997). Therefore, the net accretion rate is determined principally by the balance between productivity of the vegetation and the rate of decomposition of soil organic matter. Organic matter production and decomposition rates are determined by a myriad of factors, but those that are relevant to wetland losses in Louisiana include nutrient and sediment supply, salinity concentrations, hydroperiod, and waterlogging.

Under natural conditions, sediment deposition in wetlands is an important factor in maintaining vertical accretion by stimulating the primary production process (Hatton *et al.* 1983; Nyman *et al.* 1990; Nyman *et al.* 1993). The mineral contribution of sediment has been described as critical to increasing the elevation of the marsh, by helping maintain a surplus of production over decomposition. Thus the importance of the mineral supply is actually secondary through its contribution to soil formation by adding organic matter via stimulating net primary productivity. Activities in the Mississippi delta such as those described above have greatly reduced the level of sediment deposition in wetlands (Reed 1995). Levee construction along the Mississippi River prevents sediment deposition in coastal wetlands and this has resulted in a reduction of vertical accretion (Craig *et al.* 1979). Another contributing factor to reduced sedimentation is the extensive network of canals and impoundments in southern Louisiana. Canal spoil banks inhibit the deposition of suspended sediments and limit water exchange with


surrounding areas (Conner and Day 1987; Swenson and Turner 1987; Baumann *et al.* 1984, Boumans and Day 1990). Studies have shown that high land loss rates are directly proportional to high canal densities (Turner 1987; Turner and Cahoon 1987). Additionally, canals have contributed to the reduction of water quality by allowing nutrient-rich upland runoff to flow past wetlands directly to water bodies (Gael and Hopkinson 1979). Wetland impoundments consisting of a system of dikes and water control structures have been widely constructed in the Mississippi delta (Day *et al.* 1990). Studies have shown that these impoundments can reduce the influx of suspended sediments, lower accretion rates, lower wetland productivity, and reduce the movement of migratory marine fishes (Cahoon and Groat 1990; Cahoon 1994; Boumans and Day 1990; Flynn *et al.* 1990; and Rogers *et al.* 1993).

## 1.4 Concepts of Restoration Science

The development of ecosystem management plans to restore and rehabilitate natural resources requires an understanding of how specific ecological mechanisms regulate the structure and function of ecosystems. The idea that the specific responses of ecological systems can be projected in time under specific initial conditions is known as ‘succession’. The science of restoration ecology applies the fundamental ecological processes of succession to rehabilitate degraded landscapes, sustain technological development, and improve environmental quality (Twilley *et al.* 1998; 1999).

Restoring disturbed ecosystems to hasten their rehabilitation is simply the manipulation of ecological succession to obtain a specific goal or purpose. Knowledge of the ecological theory that pertains to ecosystem development fosters more effective restoration planning that is less expensive, can be effectively implemented, and gives a more desirable final result (Christensen *et al.* 1996). This requires diagnostic capabilities that are based on ecological theory of succession and ecosystem development. These diagnostic capabilities are presently limited by the ability of scientists to: 1) anticipate ecological responses of ecosystems to specific manipulations or site conditions; 2) monitor responses of ecosystems at sufficient space and time scales to validate these responses; and 3) modify operations of rehabilitation projects according to the response of the ecosystem to obtain specific goals. One of the most difficult tasks in restoring ecological systems is to select the proper set of criteria for site manipulations that will rehabilitate habitats to obtain a specifically defined structure and function. Thus, a fundamental need of restoration programs is to develop practical tools and approaches that can be used to predict, monitor, and validate the response of ecosystems to rehabilitation criteria.

Changes in ecosystem attributes with time, such as specific characteristics in structure or function, are known as trajectories (Figure C.1-6). Restoration ecology requires the investigation of ecological trajectories of ecosystems in response to a variety of rehabilitation conditions. In the case of coastal Louisiana, the challenge is to determine the causal linkages to the degradation of this ecosystem over the last 100 yrs (1.2 million acres), and to develop features that will move the system to some rehabilitated condition. Developing features to move a system to some desired condition of both community structure and ecosystem function requires testing hypothesis of causal mechanisms described above.



**Figure C.1-6 Changes in Ecosystem Trajectories**


The future of the Mississippi River Deltaic Plain will depend on either a strategy that will allow present rates of degradation to continue, versus mounting an effort to restore natural processes of the delta cycle (Figure C.1-7). The LCA Ecosystem Restoration Plan has adopted the term “target” to reflect the degree to which environmental drivers will be restored, and the associated ecosystem and landscape changes that would be expected over the next fifty years. Environmental drivers are the major processes external to the coast that form coastal landscapes and ecosystems (Figures C.1-4 and C.1-5). A major assumption of the LCA Restoration Plan is that interruptions of environmental drivers are responsible for the degradation of the coastal landscape by destabilizing wetland ecosystems. LCA targets are, therefore, intended to represent degrees to which important environmental drivers can be restored to stabilize wetland landscapes.

The benchmark for all restoration goals is a prediction of further landscape degradation under a ‘no action’ or ‘future without’ scenario. Again, this prediction includes the assumptions that historical causal mechanisms of ecosystem degradation, and associated rates of loss, will continue over specific time frame. Land loss rates under ‘no action’ scenario have been estimated to 2050 in the LCA plan for both the deltaic and chenier plain provinces (USGS). Restoration scenarios considered by the LCA study effort are defined by three classes: (1) reduce the present rate of wetland loss by one-half of the no-action land loss rate; (2) maintain the present total area of coastal wetlands over the next 50 yrs; (3) or enhance the present landscape condition by increasing wetland area by one-half the present annual net land loss rate over 50 yrs. These LCA restoration targets assume that a certain change in degree of environmental drivers (forcing functions in Figure C.1-5) can achieve the three restoration goals described

above (Figure C.1-7). Coastal Frameworks or features that are designed to help restore environmental drivers, such as reintroduction of the river, require estimates (projections or predictions) as to how much landscape will be built and what will be the function of resulting coastal ecosystems.

#### LCA Restoration Goals


- 1) No Action: continued degradation rate;
- 2) Reduce Loss: decrease in degradation rate;
- 3) Maintain: no wetland loss;
- 4) Enhance: increase wetland acreage.



**Figure C.1-7 Goals of the LCA Ecosystem Restoration Plan Relative to Present Assumptions of Land Loss**

The environmental drivers of a coastal landscape reflect the complex behavior of regional climate, river discharge, tides, wind, and oceanographic currents (Figure C.1-9). Coastal settings can be catalogued as river deltas, river-dominated estuaries, lagoons, and oceanic islands (Thom 1982; Woodroffe 1992). These geomorphologically distinct landforms have local variations in topography and hydrology that result in the development of particular ecological types of wetland ecosystems. Coastal wetlands ecosystems have specific community structure and function, dependent upon local effects of topography that modify the regional impacts of environmental drivers across the coastal landscape. The combination of global (climate and biogeography) and regional (geomorphology) processes modified by the local (topographic) factors determines how regulators, resources, and hydroperiod will control the patterns of wetland development (Figure C.1-8). This hierarchy in geophysical, geomorphological, and ecological processes determines the level of stress at the plant system level (core of Figure C.1-8). At any location and specific time, local factors constrain the specific attributes of the plant system in the form of gradients in resources, regulators and hydroperiod (described below). And such constraints result from basin level processes that produce a variety of subsidies and stressors to ecological processes. To achieve restoration goals, plans must effectively change

environmental drivers at the hydrologic basin level that reduce stress conditions at the local environment that are responsible for ecosystem degradation (Figure C.1-8).



Source: Gosselink as part of Coast 2050 planning document


**Figure C.1-8 Conceptual Diagram Describing the Coastal Processes Wetland Ecosystem development**

Conceptual models must be able to link processes at the province, basin, local spatial and temporal scales to recommend adjustments in environmental drivers that will allow natural processes to restore the delta and chenier plain ecosystems (Boesch *et al.* 1994). At the core of such predictions, which are based upon high scientific uncertainty, is the understanding that these assumptions represent hypotheses that must be tested during the implementation stage of the restoration program (Figure C.1-1). Determining which processes contribute significantly to the destabilization of coastal wetlands has historically been the approach of wetland ecology related to understanding impacts of land use change. Restoration objectives and goals, with the purpose of promoting marsh stabilization, represent tests of the mechanisms that were proposed above as responsible for marsh destabilization. Destabilization of marsh sediments has been described as resulting from changes in regional hydrology such as hydroperiod (Gosselink and Turner 1978; Reed and Cahoo 1992) and chronic waterlogging stress (DeLaune *et al.* 1994; Day *et al.* 1994) that are linked to reductions in plant growth (as described above). Waterlogging stress, even without added salinity stress, can be a primary cause of marsh dieback (Webb *et al.* 1995).

Succession in coastal wetlands has to account for biomass production, along with rates of soil decomposition, to understand marsh stability. Optimization theory predicts plants adapt to environmental stressors via variations in biomass allocation (Bloom *et al.* 1985; Gleeson and

Tilman 1992; Bazzaz 1997). Variations in morphology and productivity allow a plant to maximize effort in either obtaining a limited resource or responding to lethal concentrations of a plant regulator. Resources, such as nutrients and light, are defined as those elements in nature that are required for growth, and when consumed are no longer available for another individual. Regulators are defined as those physical and chemical properties that regulate physiological processes, but are not consumed by an individual, such as salinity, pH, and temperature. Stress, or reduced growth, can be manifested by the limited concentrations of resources, or by extreme concentrations of regulators. In Figure C.1-9, stress is defined as the deflection from maximum biomass levels at different concentrations of either resources (upper panel) or regulators (lower panel). The life strategy of wetland species differs from that of most terrestrial species in that tolerance to growth regulators, such as soil salinity, are as critical in plant succession as the influence of resource gradients (Grime 1977). Thus concentration gradients of both resources and regulators at the local environment (Figure C.1-8) can drive biomass allocation and succession in wetland ecosystems (Huston 1994). In addition, wetland succession is controlled directly by hydroperiod, which includes both the frequency and duration of flooding, and constitutes a third factor to consider in wetland restoration projects.

Plant productivity can be defined along these three axes that define the gradients of resources, regulators or hydroperiod as presented in the triangles in Fig. 1.9. A constraint envelope defines the productivity of coastal wetlands based on combined gradients of resources, regulators, and hydroperiod (Figure C.1-9). For most wetlands in coastal Louisiana, the most important resource gradients are bulk sediments and nutrients, regulator gradients include salinity and sulfide, and hydroperiod gradients depend on the duration of flooding. A condition exists across each gradient with maximum productivity, and this is the farthest point from the origin representing benign environment. Points close to the origin represent maximal conditions of stress. The surface area of the envelope that results from connecting the three points along the axis of each condition is proportional to net primary productivity of the wetland. The productivity of a particular wetland site can be characterized by some combination of these three conditions of the local environment, and that combination can be defined relative to the stability of a wetland. Thus, wetlands with conditions that are plotted farthest from the origin along each axis should have maximum rates of total net primary productivity compared to those plotted nearer the origin that result in severely reduced productivity.



**Figure C.1-9 Production Envelopes of Net Primary Productivity in response to the relative values of resource gradient, regulator gradient, and hydroperiod.**

The LCA Ecosystem Model uses this approach to search for optimum set of environmental drivers that will produce conditions of reduced stress and thus restore the productivity and stability of wetland ecosystems. Restoration measures should provide environmental drivers at the basin level that alter combined levels of resources, regulators, and hydroperiod at the local level that reduce the continued degradation of coastal landscape. At the same time, strategies may be necessary at the more local level that can also modify these measures at the basin level that may interfere with restoring local conditions. For example, spoil banks and canals may interfere with basin level strategies to restore resources, regulators and hydroperiod at the local environment. So that while basin strategies may be the focal point of the

LCA plan, the impact must be able to cascade from the larger to the more local level of the landscape to insure benefit to the wetland system. As the combination of resource, regulator and hydroperiod gradients changes over time, responses in primary productivity will give rise to successional patterns in wetland ecosystems. Restoration measures have to provide a combination of these three factors from the province to the local that allow optimum marsh productivity and thus sustainable coastal ecosystems.

Habitat use by fauna also follows the sequence of vegetation patterns of the delta cycle. Thus the three gradients in resources, regulators and hydroperiod can also be used to define approaches to modeling habitat suitability in the coastal landscape. The difference with higher trophic levels, compared to the focus on wetland vegetation above, is that resources are not abiotic requirements of growth (as for plants), but are spatial and organic characteristics of the landscape. Vegetation structure can be considered the structural species of an ecosystem, while most fauna represent interstitial species that colonize the physical structure provided by macrophyte vegetation (Huston 1994). Examples of interstitial species in coastal Louisiana are the marine nekton and benthic fauna that colonize wetland habitats; as well as birds that utilize vegetated areas of the landscape. This explains the general pattern that structural species have a major influence on the presence of interstitial species. Yet the use of a particular habitat depends on the spatial patchiness of emergent habitat and coastal waters – referred to as landscape ecology. This is true for both fisheries and waterfowl. In addition, models of habitat suitability must account for important regulators such as salinity, water temperature, and water flow. And as for plants, hydroperiod is important factor in habitat use.

The hierarchical approach to restoring coastal ecosystems of the Mississippi River Delta and Chenier Plain must account for the impacts of environmental drivers at the province level to ecological processes of both wetlands and waters at the local level. There are strong linkages in the landscape arrangement of geomorphic features that control not only the types of ecosystems that will develop (community structure), but also the utilization of those ecosystems by higher trophic levels (ecosystems function). The delta cycle is a complex interaction of geomorphic landscapes, geophysical processes, and ecological succession. The LCA Ecosystem Model was developed to capture the various scales by which these physical, biogeochemical, and ecological processes will change the attributes of coastal ecosystem with incremental change in environmental drivers.

## **1.5 LCA Ecosystem Model**

The goal of the Louisiana Coastal Area Restoration Plan is to rehabilitate coastal ecosystems by promoting the distribution of riverine freshwater, nutrients and sediments using natural processes that ensure the structural integrity of the estuarine basins. Developing and evaluating restoration Coastal framework features of the LCA to achieve this goal required linking the changes in environmental drivers (processes such as riverine input) to specific restoration endpoints (hydrodynamic, ecological and water quality) using a variety of modeling approaches (Figure C.1-10). This modeling effort was designed to evaluate the effects of various combinations of conceptual restoration features on the sources of ecosystem stress, identify areas of influence and project possible ecological benefits along the deltaic and chenier plains. This was accomplished by combining existing conceptual models of delta evolution and ecological succession. Assumptions of causal linkage mechanisms and desired final conditions (endpoints) were used to estimate feature requirements that will allow ecosystems to move toward target

endpoints. The endpoints were constructed into algorithms and used to calculate benefits of specific alternatives at the subprovince scale.

The linkage between restoration alternatives and restoration endpoints was provided by the construction of the LCA Ecosystem Model (Figure C.1-10). The modeling system consists of five major steps in the evaluation process. First is the development of alternatives that approximate the degree of change in environmental settings to achieve specific restoration goals (reduce, maintain, enhance, see Figure C.17). In step two, the alternatives were provided to the ecosystem modeling team for estimates of change in five different modules. These five modules included: (1) hydrodynamics, (2) land building, (3) habitat switching, (4) habitat use, and (5) water quality (Figure C.1-11). This approach is similar to the coastal ecosystem landscape models that have been developed over the last two decades to simulate processes in specific regions of coastal Louisiana (Costanza *et al.* 1988; Martin *et al.* 2000; Reyes *et al.* 2000; Martin *et al.* 2002). Each module requires knowledge of existing conditions, and will then predict changes in the landscape based on assumptions of how the ecosystems respond to coastal processes. Third, each module produced a set of endpoints specific to the environmental conditions of the particular Coastal Framework. Many of these endpoints became the input to other specific modules. The details of how these modules were linked and specifics on the modeling tools for each module are described in chapter D.2. The fourth step was to use the endpoints of these five modules in a series of benefit calculations to determine specific types of ecosystem response. Finally, the original restoration alternatives were evaluated using a collection of the benefits and compared to the original restoration objectives.




Figure C.1-10

Developing and evaluating Restoration Alternatives of Louisiana Coastal Area Restoration Plan will link the changes in forcing functions (processes such as river input) to specific restoration endpoints (hydrodynamic, ecological and water quality) using both Simulation and Desktop Modeling Approaches. Values of these endpoints and other metrics were used to calculate benefits of specific alternatives.

This chapter has described the conceptual framework of ecosystem needs in the Mississippi River Deltaic and Chenier Plain, and approach to develop a restoration plan. The trajectories described in the restoration plan and simulated (evaluated) in this appendix are based on the geophysical, geomorphic, and ecologic processes that historically formed and sustained this river-dominated coastal landscape. In the delta province, these include the formation of several deltaic lobes of the Mississippi River. In the chenier province to the west, major processes include the formation of a series of beach ridges or cheniers. These models are crude estimates as to how these linkages in coastal processes will effectively achieve a sustainable coastal landscape. Future model development will be required to evaluate the feasibility of reducing scientific uncertainty by specific changes in the design and/or operation for both existing and future projects to reach targets at the province scale. It should be understood that projections based on model development in this stage of a restoration plan are scaled to represent the basic “features” of a plan, or strategies, and not the operational nature of proposed measures.